

532014

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
6 May 2004 (06.05.2004)

PCT

(10) International Publication Number
WO 2004/037282 A1

- (51) International Patent Classification⁷: **A61K 38/44**, G01N 33/574
- (21) International Application Number: PCT/SE2003/001652
- (22) International Filing Date: 24 October 2003 (24.10.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
0203137-5 24 October 2002 (24.10.2002) SE
60/420,787 24 October 2002 (24.10.2002) US
- (71) Applicant (*for all designated States except US*):
KAROLINSKA INNOVATIONS AB [SE/SE]; S-171 77 STOCKHOLM (SE).
- (72) Inventors; and
(75) Inventors/Applicants (*for US only*): **INGEL-MAN-SUNDBERG, Magnus** [SE/SE]; Soltorpsvägen 28, S-182 33 DANDERYD (SE). **KARLGREN, Maria** [SE/SE]; Gustav III:s Boulevard 4/507, S-169 72 SOLNA (SE). **GOMEZ, Alvin** [SE/SE]; Jägargatan 20, S-118 67 STOCKHOLM (SE).
- (74) Agent: **DR LUDWIG BRANN PATENTBYRÅ AB**; Maria Skolgata 83, P.O. Box 171 92, S-104 62 STOCKHOLM (SE).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Declaration under Rule 4.17:**
— *of inventorship (Rule 4.17(iv)) for US only*
- Published:**
— *with international search report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: THE USE OF CYTOCHROME P450 ENZYME CYP2W1 AS A DRUG TRAGET FOR CANCER THERAPY

(57) Abstract: The present invention relates to a novel drug target in cancer therapy. More closely it relates to use of the cytochrome P450 enzyme CYP2W1 and its promoter as a drug target for cancer therapy. . It also relates to screening methods for obtaining therapeutic agents for cancer therapy and to therapeutic agents comprising moieties showing binding affinity for CYP2W1 and moieties with cytotoxic/anti-cancer effect.

BEST AVAILABLE COPY

WO 2004/037282 A1

DRUG TARGET IN CANCER THERAPY

Field of the invention

5 The present invention relates to a novel drug target in cancer therapy. More closely it relates to use of the cytochrome P450 enzyme CYP2W1 and its promoter as a drug target for cancer therapy. It also relates to screening methods for obtaining therapeutic agents for cancer therapy and to therapeutic agents comprising moieties showing binding affinity for CYP2W1 and moieties with cytotoxic/anti-cancer effect.

10 Background of the invention

The cytochromes P450 are responsible for the metabolism of endogenous as well as exogenous compounds. In human at present about 80 different P450 genes have been identified of which 55 are functional whereas 25 are pseudogenes. The human P450 forms can be divided into three major groups: i) those in CYP families 5-51 of endogenous importance having usually high
15 affinity for the substrates and being relatively well conserved during evolution, ii) those in CYP families 1-3, with usually less affinity for their substrates, being less conserved evolutionary and which exhibit important genetic polymorphisms and iii) those in family 4 with properties in between the two other groups (see Ingelman-Sundberg (2002) for a review). CYP enzymes in families 1-3 are active in metabolism of exogenous compounds and currently 19 genes encoding
20 active enzymes have been described. The cytochromes P450 in these families are responsible for 70-80 % of all phase I dependent metabolism of clinically used drugs and participate in the metabolism of a huge number of other xenobiotic chemicals. This can lead to the metabolic activation of precarcinogens and drugs, which might exert toxic or carcinogenic effects. This capacity to activate prodrugs to cytotoxic products are seen e.g. in the metabolic activation of
25 acetaminophen which is metabolised by CYP2E1 and CYP3A4 to a reactive imine intermediate that can cause liver hepatotoxicity. Other examples of cytochrome P450 dependent metabolic activation come from prodrugs in cancer therapy where P450 enzymes like CYP2B6 inserted by viral transfection into tumours facilitate tumour suppression upon treatment with cancer drugs

agents like e.g cyclophosphamide or ifosfamide (Waxman et al., 1999, Jounaidi and Waxman, 2001).

Summary of the invention

The invention is based on the specific expression of the cytochrome P450 enzyme CYP2W1 in transformed tissue. This tissue specific expression found by the inventors can be used to target drugs and prodrugs to the transformed cells.

The invention thus relates to a compound comprising one part conferring cytotoxic and/or anti-cancer effects to the compound and one part conferring binding affinity towards CYP2W1 to the compound. It also relates to a pharmaceutical composition comprising such a compound.

In a further aspect the invention relates to use of CYP2W1 as a drug target in both screening for therapeutic agents and in cancer therapy.

In yet a further aspect the invention relates to the CYP2W1-promoter and use thereof in screening for therapeutic agents and in gene therapy.

The invention can for example be used for, but is not restricted to, the treatment of lung tumours, colon tumours or ovarian tumours, or all or any tumour expressing CYP2W1 constitutively or after administration of an inducing agent.

Brief description of the figures

Figure 1

Gene structure (A), nucleotide and predicted amino acid sequence (B) of CYP2W1. (A) Exons are shown as boxes with size in base pairs. The size of the introns is shown in kilo base pairs. (B) The stop codon is indicated by an asterisk and the proline-rich region, PPGP, and the conserved cysteine are shown in italic bold. The 3'-flanking region was obtained from the Celera Transcript hCT1786066.

Figure 2

Unrooted phylogenetic tree of CYP2W1 and other human P450s belonging to family 1, 2 and 3. The scale bar for distance measurements represents 0.1 amino acid substitutions per site.

Figure 3

Real-time PCR based quantification of CYP2W1 mRNA expression in Human Tumour Multiple Tissue cDNA (MTC) Panel using gene specific primers.

Figure 4

5 Distribution of CYP2W1 mRNA in different human tissues and cell lines. (A) Multiple Tissue Expression (MTE) Array containing 42-500 ng of normalized human poly(A)⁺ RNA per dot. (B) Multiple Tissue Northern (MTN) blot containing approximately 1 µg poly(A)⁺ RNA per lane and Northern blot containing 20 µg total RNA from human liver, HepG2, HeLa, B16A2 or HEK293 cells per lane. All blots were hybridized with a ³²P-labeled probe corresponding to nucleotides
10 329-986 and exposed for 48 hours. The size markers used for the northern blot were the rRNA 28S (4.6-5.3 kb) and 18S (1.8-2.0 kb) band.

Figure 5

Western blot containing four lanes with HepG2 total cell extract corresponding to 40 µg protein per lane. The first two lanes were incubated with CYP2W1 antiserum, while the last two were
15 incubated with antiserum blocked with peptide. The blot was exposed for 3 minutes.

Figure 6

CO-spectral analysis of the microsomal fraction of the pCMV4-2W1 II6 construct.

Figure 7

Western blot analyses of microsomal, mitochondrial and cytosolic fractions obtained from
20 transfected HEK293 cells. Fractions corresponding to 40 µg protein were loaded per lane. The last two lanes contain human liver microsomes and cytosol corresponding to 40 µg protein. The membrane was incubated with CYP2W1 antiserum and exposed for two minutes.

Figure 8

5' flanking region deletion constructs in pGL3-Basic.

Figure 9

Comparison of luciferase activity of different promoter lengths. Luciferase assay was performed 24 hours after transfection. Basal luciferase reading (pGL3-Basic) was set at 1.0.

Figure 10

- 5 Shorter constructs made from genomic DNA. The region between the transcription initiation site and 78 bp upstream gave a significant increase in luciferase activity.

Figure 11

The region between the transcription initiation site and 78 bp upstream of it, which contains the TATA box, gives a significant increase in luciferase activity.

10

Detailed description of the invention

The approach for drug targeting in cancer therapy according to the invention is to utilize a specific form of cytochrome P450 only present in the transformed tissue being able to activate prodrugs converted to cytotoxic products. The present inventors have identified a novel human
15 cytochrome P450 form, CYP2W1, mainly expressed in tumour cells. This finding represents a major step of importance for the development of new potential agents effective in cancer therapy. Thus, in a first aspect the present invention relates to the use of the cytochrome P450 enzyme CYP2W1 and genetic variants thereof as a drug target in cancer therapy. CYP2W1 was found by the inventors to be selectively expressed in tumour tissues such as lung carcinoma, colon
20 carcinoma and ovarian carcinoma. A common problem in cancer therapy is the unselective toxicity exerted by cytotoxic drugs against cancer cells and their healthy counterparts. The present invention shows the potential to improve drug targeting to tumour cells solely by administering substances that can be metabolically activated by CYP2W1 to cytotoxic agents. Accordingly, the invention provides a method for screening for such substances, substrates for
25 CYP2W1, which can be modified by the enzyme to obtain a cytotoxic metabolite. This method comprises the steps:

1. Bringing a candidate prodrug into contact with CYP2W1 or a cell expressing CYP2W1.
2. Allowing CYP2W1 to act on the candidate prodrug to produce a candidate drug.

3. Analysing the candidate drug for cytotoxic/anti-cancer properties.

The candidate prodrug is preferably a non-toxic compound suitable for conventional pharmaceutical administration.

In another preferred method of screening for cytotoxic agents, anticancer agents, or other

5 substrates, activated by CYP2W1, the enzyme is expressed using for example yeast as expression system. Other expression systems like bacterial and mammalian expression system could also be used as enzyme source. The obtained microsomes are incubated with a fluorescent substrate that is metabolised by CYP2W1. Inhibition of that reaction is then used for drug screening with different substrates. Candidate substrates to be tested, but not restricted to, could be drugs with
10 moieties that are known to be metabolised by other members of the CYP2 family. The substrate that shows the best inhibition can be modified in order to obtain a cytotoxic metabolite. Thus the invention relates to a screening assay using CYP2W1 as a drug target.

Another aspect of the invention relates to targeting of drugs and prodrugs to cancer cells,

15 preferably to lung, colon or ovarian tumour cells. Since cytochromes P450 are partially expressed on the cell surface, drugs and/or prodrugs can also be targeted to cancer cells by the coupling of them to ligands which bind specifically to CYP2W1. In this aspect, "prodrug" means a substance which will be converted to a cytotoxic and/or anti-cancer drug upon binding to a cell expressing CYP2W1 on its surface. The drug to be administered in this way could be any

20 anticancer/cytotoxic drug known in the art. The prodrug could be a prodrug identified by the screening method described above or any other compound which is activated to a cytotoxic or anti-cancer agent by CYP2W1. The ligand to which the drug/prodrug should be coupled could be any molecule that binds specifically to CYP2W1 bound on the cell surface, i.e. does not show any substantial specific binding affinity towards any other endogenous or exogenous substances.

25 The ligand could be an antibody, preferably a monoclonal antibody or a binding fragment thereof, which can be prepared by standard techniques. Pharmaceutical compositions comprising the drug/prodrug-ligand compound could be prepared by techniques known in the art.

Yet another aspect of the invention relates to the combinatorial use of a prodrug to be

30 metabolised, and a substance that can induce the expression of CYP2W1 in the tumour cells, thus resulting in higher concentrations of the cytotoxic metabolite at the site of the tumour .

In a further aspect, the invention relates to a method of providing therapeutic agents for cancer therapy, comprising screening for such agents by using CYP2W1 as a drug target.

The therapeutic agents shall modulate the activity of CYP2W1, such as induce or increase the activity thereof. This method comprises the steps:

1. Bringing a candidate agent into contact with a cell capable of expressing CYP2W1 or a reporter gene operably linked to the regulatory sequences of CYP2W1.
2. Measuring changes in expression and/or activity of CYP2W1

In order to screen for agents inducing the expression of CYP2W1 different cell lines can be used.

One example of possible cell line is the human hepatoma cell line HepG2, which is constitutively expressing CYP2W1. The agent to be tested will be administered to the supplement medium and the induction of the CYP2W1 expression can be tested on the RNA, protein and/or enzymatic activity levels. Candidate agents to be tested, but not restricted to, could be known inducers of cytochrome P450s belonging to the CYP2 and CYP3 family, including those being efficient substrates for the nuclear receptors PXR, LXR, FXR and CAR.

In a yet further aspect, the invention relates to a method of treating cancer, comprising giving a patient in need thereof a therapeutically effective amount of a substance activated by the enzyme CYP2W1 and/or inducing the enzyme CYP2W1.

For example, the substance may be activated to an anticancer agent or a cytotoxic substance.

The invention also relates to the use of the CYP2W1 promoter (SEQ ID NO: 10) in gene therapy. This aspect includes the use of the promoter in the manufacture of a medicament, preferably a medicament for use in cancer therapy, even more preferably for use in treatment of lung tumours, colon tumours or ovarian tumours. This aspect also includes the use of DNA-molecules containing the CYP2W1 promoter in pharmaceutical compositions, such DNA-molecules could be, for example, vectors for delivery of the promoter to the patient in need thereof.

In an even further aspect the invention relates to the use of the CYP2W1 promoter in screening for modulators, such as inducers or enhancers, of CYP2W1. The promoter can be linked to a variety of reporter systems known in the art in order to study the effects of candidate modulators.

Experimental

The present invention will be described in more detail below in an experimental section.

Materials and Methods**5 Bioinformatics**

The BLASTN search algorithm was used to search the Celera sequence databases (<http://www.celera.com>) for sequences related to the previously reported CYP2W1 partial cDNA sequence (GenBank Accession No. AK000366). The amino acid sequence of CYP2W1 was compared with the amino acid sequences of other human P450s using multiple sequence
10 alignment. An unrooted phylogenetic tree was calculated using ClustalW 1.8 (Thompson et al., 1997) and visualized using the program unrooted (<http://pbil.univ-lyon1.fr/software/unrooted.html>).

RNA Isolation and 1st strand cDNA Synthesis

Total RNA was isolated from cultured HepG2 and B16A2 cells using the TRIZOLE Reagent
15 (Invitrogen, Rockville, MD) based on the acid guanidine thiocyanate phenol chloroform extraction method, whereas total RNA from HEK293 and HeLa cells was isolated using the RNeasy Mini Kit (Qiagen, Hilden, Germany) Reverse transcriptase was performed using MMLV Reverse Transcriptase (Invitrogen, Rockville, MD) according to the manufacturer's instructions.

Amplification and Sequencing

20 cDNA from HepG2 cells was used as a template in PCR reactions with CYP2W1 specific primers. The enzyme used for amplification was Taq DNA Polymerase (ABgene, Epsom, UK). The PCR products generated were sequenced using the ABI PRISM BigDye Terminator Cycle Sequencing Ready Reaction kit (Applied Biosystems, Foster City, CA), according to the manufacturers instructions, and analyzed on an ABI PRISM 377 DNA Sequencer (Applied
25 Biosystems, Foster City, CA).

Multiple Tissue cDNA Panels

cDNA from Human Multiple Tissue cDNA (MTC) Panel I and II, Human Digestive System MTC Panel, Human Fetal MTC Panel and Human Tumour MTC Panel (Clontech, Palo Alto,

CA) were used as templates in a PCR-reaction using the CYP2W1 specific primer pair, 5'-GAGGTGGAGGCATCTTCTTCTCATCTG-3'/5'-CTGGATCAGGGCGTCCACATAGCTG-3'. A reaction without cDNA was run as a negative control in each experiment. The PCR amplification was performed in the presence of 1 mM MgCl₂, 0.2 mM of each dNTP, 0.4 µM of each primer, 0.625 U Taq DNA Polymerase (ABgene, Epsom, UK), 1x Reaction Buffer IV (75 mM Tris-HCl pH 8.8 at 25°C, 20 mM (NH₄)₂SO₄, 0.01% (v/v) Tween 20) and 0.5 ng cDNA. The PCR reaction was performed by an initial denaturation step at 94°C for 3 min, after which the amplification was carried out for the indicated number of cycles, with denaturation at 94°C for 45s, annealing at 60°C for 30s, and extension at 72°C for 1 min and 30s, and a final extension step of 72°C for 10 min. HepG2 cDNA was used as a positive control, and the quality of the cDNA in the MTC panels was ensured by using the G3PDH control primers, supplied by the manufacturer. Amplification was performed under the same conditions as described above. The PCR products were separated and detected on a 1.2% ethidium-bromide stained agarose gel. The PCR product generated was sequenced and analyzed, as described above, in order to ensure specificity of the reaction.

Real-time quantitative PCR detection

CYP2W1 mRNA expression was quantified in Human Tumour MTC Panel using the Smart Cycler® System (Cepheid, Sunnyvale, CA). The quantitative real-time PCR assay was carried out using the CYP2W1 gene specific primer pair, 5'-AGCTATGTGGACGCCCTGATCCA-3'/5'-ACGCGGTCTAGCTCCTCCTGCAC-3', and SYBR®Green PCR Master Mix (Applied Biosystems, Foster City, CA). The quantification was performed using 0.5 ng of each cDNA sample. To avoid detection of any possible contaminating genomic DNA the primer pair was designed so the resulting product spans exon junctions. Standard curves were constructed with the use of serial 10-fold dilutions ranging from 10⁻² fmol/µl to 10⁻⁹ fmol/µl of an accurately determined concentration of a CYP2W1 full-length cDNA fragment. The data was analyzed using the Smart Cycler® Software (Cepheid, Sunnyvale, CA).

Analysis of mRNA by Northern and Dot Blot

A Human Multiple Tissue Expression (MTE) Array (Clontech, Palo Alto, CA) and a Multiple Tissue Northern (MTN) Blot (Clontech, Palo Alto, CA) were hybridized with a ³²P-labeled

CYP2W1 probe, corresponding to nucleotides 329-986, which was generated by PCR and labeled using the Radprime DNA labelling system (Invitrogen, Rockville, MD). Hybridization was performed according to the manufacturer's instructions.

The ³²P-labeled CYP2W1 probe was also hybridized to total RNA from HepG2, B16A2,

5 HEK293 and HeLa cells and total RNA from human liver (provided by Anna Westlind). In particular, 20 µg RNA per lane was loaded on a 1.2% agarose/formaldehyde denaturing gel and subjected to electrophoresis using standard procedures (Struhl, 1993). The RNA was transferred to a Hybond-N+ filter (Amersham Pharmacia Biotech, Uppsala, Sweden) and hybridized using ExpressHyb Hybridization Solution (Clontech, Palo Alto, CA). Prehybridization was carried out
10 at 65°C for 30 minutes, and hybridisation was carried out at the same temperature over night.

After hybridisation all three blots were exposed to an imaging plate and analysed by a Bas-1800II phosphoimager using the Image Gauge software (Fujifilm, Samford, CT).

Cell Culture Conditions

HepG2 and HEK293 cells were cultured in Minimum Essential Medium supplemented with 10%
15 fetal bovine serum, 1mM sodium pyruvate, 1% non-essential amino acids, 100 IU/ml penicillin and 100 µg/ml streptomycin. HeLa cells were cultured in Dulbecco's Modified Eagle Medium supplemented with 10% fetal bovine serum, 1% non-essential amino acids, 100 IU/ml penicillin and 100 µg/ml streptomycin. B16A2 cells were cultured in William's E Medium supplemented with 5% fetal bovine serum, 0.6 µM hydrocortisone, 0.25 µg/ml insulin, 100 IU/ml penicillin and
20 100 µg/ml streptomycin. All cells were cultured in humidified 5% CO₂ at 37°C. All cell culture media and supplements were obtained from Invitrogen, Rockville, MD.

CYP2W1 Antiserum

An 15 amino acid long peptide, TMRPRAQALCAVPRP, corresponding to the C-terminus of CYP2W1 was synthesized and coupled to keyhole limpet hemocyanin (Neosystems, Strasbourg,
25 France). Immunization of two rabbits was carried out with injections at day 0, 14, 27 and 56. Bleedings were performed before the first injection to obtain pre-immune serum, and after 39, 67 and 96 days to obtain the CYP2W1-specific antiserum.

Preparation of Total Cell Extracts and Western Blot

The human cell lines HepG2, HEK293, HeLa and B16A2 were analysed for CYP2W1 expression using western blot. Cells were washed twice, scraped into ice-cold PBS and pelleted by centrifugation at 17,320x g. The pelleted cells were resuspended in a phosphate buffer, containing 50 mM NaPO₄ pH 7.4, 0.1mM EDTA and 10% glycerol, and lysed by sonication. After centrifugation at 17,320x g, supernatants were collected and protein concentration was determined according to Lowry. 40 µg total cell extract as well as human liver microsomes and cytosol, were subjected to SDS-PAGE using a 10% gel. The proteins were subsequently transferred to a Hybond-C extra membrane (Amersham Pharmacia Biotech, Uppsala, Sweden). After transfer, the membrane was blocked in TBS containing 0.05% (v/v) Tween 20 and 5% fat-free milk, and incubated with a 1:5000 dilution of the CYP2W1 antiserum. This was followed by an incubation with 1:2000 diluted horseradish peroxidase-conjugated anti-rabbit immunoglobulins and detection using SuperSignal West Pico Chemiluminiscent Substrate (Pierce, Rockford, IL). Blocking experiments were performed using antiserum pre-incubated with peptide. The CYP2W1 antiserum was incubated together with the peptide used for immunisation, at a concentration of 3 µg peptide per µl antiserum. Incubation was performed at room temperature for 2 hours under gentle agitation. After incubation, the antiserum was diluted and immunoblotting was carried out, as described above.

Construction of CYP2W1 Expression Plasmids

A full-length cDNA clone of CYP2W1 was amplified from cDNA derived from HepG2 cells using Elongase Enzyme Mix (Invitrogen, Rockville, MD). The primer pair used was 5'-GACAGATCTATGGCCCTGCTGCTCTTG-3'/5'-GACTCTAGACTAGGGCCTGGGCACCGCA-3'. Restriction sites for *Bgl*II and *Xba*I (underlined) were introduced by the primers. The PCR reaction contained 0.2 mM of each dNTP, 0.4 µM of each primer, 2.5 U Elongase Enzyme Mix (Clontech, Palo Alto, CA), and a Buffer A/Buffer B ratio corresponding to 1.7 mM MgCl₂ (5x Buffer A containing 300 mM Tris-SO₄ pH 9.1, 90 mM (NH₄)₂SO₄, and 5 mM MgSO₄; 5x Buffer B containing 300 mM Tris-SO₄ pH 9.1, 90 mM (NH₄)₂SO₄, and 10 mM MgSO₄). The conditions for the reaction were as follows, a denaturation step at 94°C for 1 min followed by 70°C for 1 min when Buffer A/Buffer B and Elongase Enzyme Mix were added as a hot start. Before amplification an additional denaturation

step at 94°C for 1 min was performed. The amplification was carried out with denaturation at 94°C for 30s, annealing at 52°C for 30s, and extension at 68°C for 2 min for 35 cycles, and a final extension step of 68°C for 7 min. The obtained PCR product was digested with *Bgl*II and *Xba*I (Invitrogen, Rockville, MD) and ligated into the expression vector pCMV4, cut with the same restriction enzymes as the PCR product. The sequence inserted was verified by DNA sequencing.

Heterologous Expression in Mammalian Cells

The HEK293 cell line was used for expression of the CYP2W1 protein. The cells were transfected with 30 µg pCMV4-2W1 plasmid per 15 cm cell culture dish using Lipofectamine 2000 (Invitrogen, Rockville, MD) according to the manufacturers instructions. Cells transfected with only pCMV4 was used as a negative control. The cells were harvested 66 h after transfection and homogenized in 10 mM Tris/HCl pH 7.4 containing 1 mM EDTA, 0.5 M sucrose and Complete protease inhibitor (Roche, Mannheim, Germany). The homogenate was fractionated by centrifugation, and the microsomal fraction was used for estimation of the total P450 content. The microsomal, mitochondrial and cytosolic fractions were subjected to western blot for detection of the CYP2W1 protein, as described above.

Identification of promoter

Varying lengths of the 5' flanking region of the *CYP2W1* gene were subcloned into the reporter vector, pGL3-Basic. These were transfected into the hepatoma cell line HepG2 (Lipofectamine 2000, Invitrogen, and Tfx-20 Reagent, Promega) 24 hours prior to luciferase assay (Dual-Luciferase Reporter Assay System, Promega). The region that gave the highest luciferase activity was further analyzed by preparing constructs of shorter intervals. A transcription factor search was the basis for making these constructs.

Results

A number of genomic clones related to CYP2W1 were identified and aligned together in order to obtain the *CYP2W1* gene sequence. A Celera transcript, hCT1786066, was also identified which

contained the 3' untranslated region of the cDNA. The *CYP2W1* gene was more than 5.5 kb long and showed the typical family 2 gene structure with nine exons (Fig. 1A). The obtained gene sequence was used when designing primers for the amplification and sequencing of the entire HepG2 CYP2W1 cDNA. Sequencing resulted in an open reading frame of 1473 nucleotides, which encodes a 490 amino acid long polypeptide. The nucleotide and predicted amino acid sequence of CYP2W1 are shown in figure 1B. CYP2W1 was found to contain some typical structural features associated with P450s, including a hydrophobic NH₂-terminal, the proline-rich region and the conserved cysteine, which is the fifth heme iron ligand. The CYP2W1 cDNA sequence described here differs extensively from the previously reported partial sequence (GenBank Accession No. AK000366) with respect to the N- and C-terminal exons.

An unrooted phylogenetic tree of CYP2W1 and other human P450s is shown in figure 2. It shows that CYP2W1 align together with the other family 2 members, but it does not seem to be closely related to any of them. The enzymes within family 2 with highest identity to CYP2W1 are CYP2D6 (42%) and CYP2S1 (40%).

The tissue-specific CYP2W1 mRNA distribution was analysed by using MTC panels by PCR. The cDNA samples in the MTC panels were pooled from several individuals and normalized with at least four different housekeeping genes. As shown in Fig 3 and Table 1 the tissues that showed the highest expression of CYP2W1 mRNA were the tumour tissues lung carcinoma and colon adenocarcinoma. Moderate expression was seen in for example fetal lung, but the expression levels seen in the majority of normal adult and fetal tissues were low or absent. A rough calculation about the difference in expression between the tumour cells having highest CYP2W1 mRNA and normal tissue reveals a 10,000-100,000 fold difference.

Table 1

Tissue specific expression of CYP2W1 revealed by PCR analyses using gene specific primers and different Human Multiple Tissue cDNA (MTC) Panels. The number of cycles required to obtain a visible product band on an ethidium-bromide stained gel is given in the column. The abbreviation nd stands for not detected, which means that no product band could be seen after 40 cycles of amplification.

Panel	Tissue	Cycles for Detection
Human MTC Panel I	Brain	nd
	Heart	nd
	Kidney	36-40
	Liver	36-40
	Lung	nd
	Pancreas	31-35
	Placenta	36-40
	Skeletal Muscle	36-40
Human MTC Panel II	Colon	31-35
	Ovary	36-40
	Peripheral Blood Leukocytes	nd
	Prostate	31-35
	Small Intestine	nd
	Spleen	nd
	Testis	nd
	Thymus	nd
Human Digestive System MTC Panel	Cecum	36-40
	Colon, Ascending	36-40
	Colon, Descending	36-40
	Colon, Transverse	36-40
	Duodenum	31-35
	Esophagus	36-40
	Ileocecum	36-40
	Ileum	31-35
	Jejunum	26-30
	Liver	31-35
	Rectum	26-30
	Stomach	31-35

Human Fetal MTC Panel	Brain	nd
	Heart	nd
	Kidney	31-35
	Liver	31-35
	Lung	26-30
	Skeletal Muscle	nd
	Spleen	nd
	Thymus	nd
Human Tumour MTC Panel	Breast Carcinoma (GI-101)	nd
	Lung Carcinoma (LX-1)	21-25
	Colon Adenocarcinoma (CX-1)	21-25
	Lung (GI-117)	36-40
	Prostatic Adenocarcinoma (PC3)	31-35
	Colon Adenocarcinoma (GI-112)	36-40
	Ovarian Carcinoma (GI-102)	26-30
	Pancreatic Adenocarcinoma (GI-103)	nd

The mRNA distribution was also examined using different mRNA blots, like MTE Array, MTN blot and Northern blotting. The mRNA amounts on the MTE Array have been normalized using eight different housekeeping genes, and the MTN blot has been normalized with respect to β -actin. Using these two methods for mRNA analysis no signals, and hence no expression, could be observed in any of the tissues (Fig. 4A and B). A Northern blot analysis was also performed using RNA from HepG2, B16A2, HEK293 and HeLa cells and human liver. In HepG2 cells a strong signal corresponding to a transcript of approximately 2.3 kb was observed (Fig.4B), which corresponds well to the sequence shown in Figure 1B. No signals could be obtained in the other cell lines or in human liver, which is consistent with the low CYP2W1 mRNA expression seen in liver using the MTC panels.

A CYP2W1 antiserum was produced by immunization of rabbits, using the C-terminal sequence of CYP2W1. The cell lines examined for the presence of CYP2W1 using western blotting were HepG2, HEK293, HeLa and B16A2, and in addition, we also determined the content in human liver microsomes and cytosol. The membranes containing HepG2 total cell extract and incubated with CYP2W1 antiserum recognized two bands of approximately 54 and 52 kDa as seen in

Figure 5. The upper band is the result of unspecific binding to a cytosolic protein, whereas the size of the lower band corresponds well to the calculated CYP2W1 protein sequence of 490 amino acids. When immunoblotting was made using pre-immune serum (data not shown) or antiserum blocked with the immunizing peptide the previously seen bands were abolished (Fig. 5). No band corresponding to CYP2W1 was seen in human liver or in the other cell lines examined (data not shown). Cross-reactivity with other P450s could be excluded since no band was observed in human liver microsomes, which normally shows high expression of most P450s in families 1-3.

CYP2W1 was heterologously expressed in HEK293 cells using three different pCMV4-2W1 constructs (II6, III6 and III7) and empty pCMV4 vector as a negative control. The microsomal fraction of the transfected cells was used for an estimation of the P450 content by spectral analysis of the reduced CO-bound form (Fig. 6). A peak was observed at approximately 450 nm indicating presence of active P450 enzyme. The amount of P450 obtained was 5 pmol/mg microsomal protein. The microsomal, mitochondrial and cytosolic fractions were subjected to western blotting for detection of the CYP2W1 protein (Fig. 7). A strong band indicating CYP2W1 were seen in the microsomal and mitochondrial fractions transfected with all pCMV4-2W1 constructs whereas no bands could be seen in the negative control.

Different parts of the 5'-flanking region of the CYP2W1 gene were subcloned to a reporter vector pGL3-Basic which bears the luciferase gene. Hepatoma cells HepG2 were used for the transfection experiments. Initial analysis of different lengths of the *CYP2W1* promoter region (Figure 8) suggests that the region between 130 bp and 500 bp upstream of the transcription initiation site (Figure 9, Subject 212) may possess a binding site for a transcription factor that can maximally drive the expression of CYP2W1. Shorter fragments from this region were analyzed (Figure 10). The shortest fragment included in the experiment gave a significant increase in luciferase activity (Figure 10) and this region contains the TATA box where RNA polymerase II binds and initiates transcription (Figure 11). Several transcriptional factors and activators are expected to bind to this region and may also affect the level of expression of CYP2W1.

It is concluded that the region between -1 and -130 bp, SEQ ID NO: 10, constitutes a very strong promoter for expression of CYP2W1 in the hepatoma cells.

Discussion

5 Alignment of genomic clones allowed the construction of the gene structure of CYP2W1. Based upon this it was possible to make a full length cDNA construct using mRNA from HepG2 cells which revealed a P450 enzyme with expected length and sequence motifs. Semiquantitative analysis for the amount of mRNA expressed in human MTC panels revealed the highest expression in a lung, a colon and an ovarian tumour, whereas in general only small mRNA
10 expression was seen in preparations from adult tissues. Based upon the number of cycles during PCR necessary to amplify the product a rough calculation reveals 10,000-100,000 fold higher CYP2W1 mRNA expressed in these tumour cells than in non-transformed cells. Further quantitative estimations are now done using Northern blot filters from different human tumours. The inventors detected significant mRNA expression in HepG2 cells whereas, by contrast, they
15 could not see any significant expression as determined by Northern blotting in different tissues from adult human.

The inventors successfully obtained a specific antiserum to the protein. Western blotting experiments revealed that no significant expression was seen in human liver, but that high
20 expression was seen in the transformed cell line HepG2. This is promising in order to obtain specific Western blotting analyses for detection of the enzyme in different tumours. In HEK293 cells heterologously expressed with CYP2W1 the expression of three bands was seen with slightly different molecular weights. The significance of this observation is currently evaluated.

25 Large cultivation of HEK293 cells transfected with pCMV-CYP2W1 cDNA allowed the preparation of microsomes useful for detection of spectrally identifiable CYP2W1. The results showed a typical P450 spectrum indicating proper folding of the enzyme being catalytically active.

30 The transformed cells line HepG2 has not been described to express any known form of microsomal cytochrome P450 at significant levels. Unexpectedly, the present inventors found

relatively high CYP2W1 expression as revealed by the Western blotting experiments in HepG2 cells. The background for this is unknown but current work aims at identifying the transcription factors and cis acting elements required for this expression. In addition, the semiquantitative determination of CYP2W1 in some different human tumours revealed an exceptionally high expression.

The present inventors propose use of the enzyme CYP2W1 as a drug target for the development of cytotoxic drugs aimed for cancer therapy. Previously the extrahepatic CYP1B1 form has been shown to be more abundantly expressed in tumours than in non-transformed tissue whereas CYP2W1 appears to represent a much more important example of tumour specific cytochrome P450 expression.

References

Ingelman-Sundberg M., Cytochrome P450 enzymes,. *Naunyn-Schmiedeberg's archives of pharmacology*, 2003, in press.

Jounaidi Y, Waxman DJ. Frequent, moderate-dose cyclophosphamide administration improves the efficacy of cytochrome P-450/cytochrome P-450 reductase-based cancer gene therapy. *Cancer Res.* 2001 ;61:4437-44.

Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG. The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* 1997 ; 25:4876-82.

Waxman DJ, Chen L, Hecht JE, Jounaidi Y. Cytochrome P450-based cancer gene therapy: recent advances and future prospects. *Drug Metab Rev.* 1999;31:503-22.

CLAIMS

1. A compound comprising one part conferring cytotoxic and/or anti-cancer effects to the compound and one part conferring binding affinity towards CYP2W1 to the compound.
5
2. A pharmaceutical composition comprising a compound according to claim 1 and pharmaceutically acceptable excipients and/or carriers.
3. An antibody, preferably a monoclonal antibody, binding specifically to CYP2W1.
10
4. Use of the cytochrome P450 enzyme CYP2W1 and genetic variants thereof as a drug target in cancer therapy, preferably in the treatment of lung tumours, colon tumours and/or ovarian tumours.
- 15 5. A method of providing therapeutic agents for cancer therapy, comprising screening for such agents by using CYP2W1 as a drug target.
6. A method according to claim 5, comprising screening for therapeutic agents modulating, preferably increasing, the activity of CYP2W1.
20
7. A method of treating cancer, comprising administering to a subject in need thereof a therapeutically effective amount of a substance activated by the enzyme CYP2W1 and/or inducing the enzyme CYP2W1 and/or the compound according to claim 1.
- 25 8. DNA-molecule with nucleotide sequence according to SEQ ID NO: 10.
9. Use of a DNA-molecule according to claim 8 in the manufacture of a medicament.
10. A method of providing therapeutic agents for cancer therapy, comprising screening for such agents modulating expression of genes regulated by the CYP2W1 promoter according to SEQ
30 ID NO: 10.

Fig 1A 1/15




Fig 1B

2/15

atg gcc ctg ctg ctc ttg ctg ttc ctg ggc ctc ctg ggg ctc tgg ggg	48
Met Ala Leu Leu Leu Leu Leu Phe Leu Gly Leu Leu Gly Leu Trp Gly	
1 5 10 15	
ctg ctc tgc gcc tgc gcc caa gac ccc tcc cca gct gcc cgg tgg ccc	96
Leu Leu Cys Ala Cys Ala Gln Asp Pro Ser Pro Ala Ala Arg Trp Pro	
20 25 30	
ccg ggg cct cgc ccg ctg ccg ctc gtc ggg aac ctg cac ttg ctg cgt	144
Pro Gly Pro Arg Pro Leu Pro Leu Val Gly Asn Leu His Leu Leu Arg	
35 40 45	
ctg tgc caa cag gac cgg tcc ctg atg gag ctc tca gaa cgc tac ggg	192
Leu Ser Gln Gln Asp Arg Ser Leu Met Glu Leu Ser Glu Arg Tyr Gly	
50 55 60	
ccg gtg ttc acc gtg cac ctg ggg cgc cag aag acg gtg gtg ctg acg	240
Pro Val Phe Thr Val His Leu Gly Arg Gln Lys Thr Val Val Leu Thr	
65 70 75 80	
ggg ttc gag gcg gtc aaa gag gcg ctg gcg ggc ccc ggg cag gag ctg	288
Gly Phe Glu Ala Val Lys Glu Ala Leu Ala Gly Pro Gly Gln Glu Leu	
85 90 95	
gcc gac cgg cct ccc atc gcc atc ttc cag ctc atc cag cga ggt gga	336
Ala Asp Arg Pro Pro Ile Ala Ile Phe Gln Leu Ile Gln Arg Gly Gly	
100 105 110	
ggc atc ttc ttc tca tct ggg gcg cgc tgg agg gct gcc cgc cag ttc	384
Gly Ile Phe Phe Ser Ser Gly Ala Arg Trp Arg Ala Ala Arg Gln Phe	
115 120 125	
acg gtg cgt gcc ctg cac agc ctg ggc gtg ggc cgg gag ccg gtg gct	432
Thr Val Arg Ala Leu His Ser Leu Gly Val Gly Arg Glu Pro Val Ala	
130 135 140	
gac aag att ctg cag gag ctg aaa tgc ctc tct ggg cag ctg gat ggc	480
Asp Lys Ile Leu Gln Glu Leu Lys Cys Leu Ser Gly Gln Leu Asp Gly	
145 150 155 160	
tac aga ggc cgg ccc ttc ccg ctg gcc cta ctg ggc tgg gct ccc tcc	528
Tyr Arg Gly Arg Pro Phe Pro Leu Ala Leu Leu Gly Trp Ala Pro Ser	
165 170 175	
aat atc acc ttc gcg ctc ctc ttc ggc cgc cga ttt gac tac cgg gac	576
Asn Ile Thr Phe Ala Leu Leu Phe Gly Arg Arg Phe Asp Tyr Arg Asp	
180 185 190	
ccc gtg ttt gtg tcc ctg ctg ggt ctc atc gat gag gtc atg gtc ctc	624
Pro Val Phe Val Ser Leu Leu Gly Leu Ile Asp Glu Val Met Val Leu	
195 200 205	

Fig 1B (cont.)

3/15

ttg ggg tcc cct ggc ctg cag ctg ttc aac gtc tac cca tgg ctc ggg Leu Gly Ser Pro Gly Leu Gln Leu Phe Asn Val Tyr Pro Trp Leu Gly 210 215 220	672
gcc ctg ctc cag ctg cac cgg ccc gtc ctg cgc aag atc gag gag gtc Ala Leu Leu Gln Leu His Arg Pro Val Leu Arg Lys Ile Glu Glu Val 225 230 235 240	720
cgt gcc att ctg agg acc ctc ctg gag gcg cgg agg ccc cac gtg tgc Arg Ala Ile Leu Arg Thr Leu Leu Glu Ala Arg Arg Pro His Val Cys 245 250 255	768
ccg ggg gac ccc gtg tgc agc tat gtg gac gcc ctg atc cag cag gga Pro Gly Asp Pro Val Cys Ser Tyr Val Asp Ala Leu Ile Gln Gln Gly 260 265 270	816
cag ggg gat gac ccc gag ggc ctg ttt gct gag gcc aac gcg gtg gcc Gln Gly Asp Asp Pro Glu Gly Leu Phe Ala Glu Ala Asn Ala Val Ala 275 280 285	864
tgc acc ctg gac atg gtc atg gcc ggg acg gag acg acc tcg gcc acg Cys Thr Leu Asp Met Val Met Ala Gly Thr Glu Thr Thr Ser Ala Thr 290 295 300	912
ctg cag tgg gcc gca ctt ctg atg ggc cgg cac ccg gac gtg cag ggc Leu Gln Trp Ala Ala Leu Leu Met Gly Arg His Pro Asp Val Gln Gly 305 310 315 320	960
cgg gtg cag gag gag cta gac cgc gtg ctg ggc cct ggg cgg act ccc Arg Val Gln Glu Glu Leu Asp Arg Val Leu Gly Pro Gly Arg Thr Pro 325 330 335	1008
cgg ctg gag gac cag cag gct ctg ccc tac aca agc gcc gtg ctc cac Arg Leu Glu Asp Gln Gln Ala Leu Pro Tyr Thr Ser Ala Val Leu His 340 345 350	1056
gag gtg cag cgg ttc atc acg ctc ctg ccg cac gtg ccc cgc tgc acc Glu Val Gln Arg Phe Ile Thr Leu Leu Pro His Val Pro Arg Cys Thr 355 360 365	1104
gcg gcc gac aca cag ctg ggc ggc ttc ctg ctc ccc aag ggc acg ccc Ala Ala Asp Thr Gln Leu Gly Gly Phe Leu Leu Pro Lys Gly Thr Pro 370 375 380	1152
gtg att ccc ctg ctg acc tcg gtg ctc ctg gat gag aca cag tgg cag Val Ile Pro Leu Leu Thr Ser Val Leu Leu Asp Glu Thr Gln Trp Gln 385 390 395 400	1200
acc cca ggc cag ttc aac ccc ggc cat ttc ctg gac gcg aat ggg cac Thr Pro Gly Gln Phe Asn Pro Gly His Phe Leu Asp Ala Asn Gly His 405 410 415	1248

Fig 1B (cont.)

4/15

ttt gtg aag cgg gag gcc ttc ctg cct ttc tct gca ggc cgc cgc gtc	1296
Phe Val Lys Arg Glu Ala Phe Leu Pro Phe Ser Ala Gly Arg Arg Val	
420 425 430	
tgt gtt ggg gag cgc ctg gcc agg acc gag ctc ttc ctg ctg ttt gcc	1344
Cys Val Gly Glu Arg Leu Ala Arg Thr Glu Leu Phe Leu Leu Phe Ala	
435 440 445	
ggc ctc ctg cag agg tac cgc ctg ctg ccc ccg cct ggc gtc agt ccg	1392
Gly Leu Leu Gln Arg Tyr Arg Leu Leu Pro Pro Pro Gly Val Ser Pro	
450 455 460	
gcc tcc ctg gac acc acg ccc gcc cgg gct ttt acc atg agg ccg agg	1440
Ala Ser Leu Asp Thr Thr Pro Ala Arg Ala Phe Thr Met Arg Pro Arg	
465 470 475 480	
gcc cag gcc ctg tgt gcg gtg ccc agg ccc taggagctcc cccagcccc	1490
Ala Gln Ala Leu Cys Ala Val Pro Arg Pro	
485 490	
aggtcctcct gaccactccc ctcccagccc tgggtcctcc caccctctct cctcccaccc	1550
cacagctcgg actgctctgg gagggccctg aggactccca ccctcacccc cacccccaca	1610
gggtcagcaa ctgcttccgg ttacaccag gactaccct gcccgaccct gtgggacccc	1670
caccctctg atgctgtctg cagctcagtc cctgccagcc ccagggagcg cctccagggc	1730
cccgccact ctcccacccc tgaagctgca ctcccaccca cctagctccc cccagggccc	1790
cccagcacct acagctgggg ctgcagggag acaacgggtg gctgcatcca gccagagaca	1850
ggcgcaggtg ggtgtcctca gcgtgcgagc cctgcacccc ccaggtcctg ggactcctgc	1910
agaccccact ccattcccgc tcctggaaca cttcctgcag ctgtgcctgg aggcagtcgg	1970
cctgcagtgc cagactctga gccaaaccac tggggccatg cgtatgactg gtgcagggag	2030
gcaaggccca cattctcctt cagagacagg cactggcgcc agaggcttcc ttggggcggg	2090
gggagggcac ctccagcccct gaagacaagc agcactgcag tggcaaaaat ggaaacactg	2150
accgggtgcg gtggctcatg cctgtaatcc cagcactttg ggaggccgag gcaggcgaat	2210
cacgaggtca ggagttcgag accagcctgc ccaacatggt gaaaccctgt ctctactaaa	2270
aatacaaaaa aattaaccga gcatg	2295

Fig 2 5/15 0.1




Fig 3 6/15





Fig 4A 7/15


	1	2	3	4	5	6	7	8	9	10	11	12
A	whole brain	cerebellum left		heart	esophagus	colon transverse	kidney	lung	liver	leukemia HL-60	fetal brain	yeast total RNA
B	cerebral cortex	cerebellum right	accumbens nucleus	aorta	stomach	colon descending	skeletal muscle	placenta	pancreas	HeLa S3	fetal heart	yeast tRNA
C	frontal lobe	corpus callosum	thalamus	atrium left	duodenum	rectum	spleen	bladder	adrenal gland	leukemia K-562	fetal kidney	<i>E. coli</i> rRNA
D	parietal lobe	amygdala		atrium right	jejunum		thymus	uterus	thyroid gland	leukemia MOLT-4	fetal liver	<i>E. coli</i> DNA
E	occipital lobe	caudate nucleus		ventricle left	ileum		peripheral blood leukocyte	prostate	salivary gland	Burkitt's lymphoma Raji	fetal spleen	Poly r(A)
F	temporal lobe	hippocampus		ventricle right	ileocecum		lymph node	testis		Burkitt's lymphoma Daudi	fetal thymus	human C ₀ t-1 DNA
G	paracentral gyrus of cerebral cortex	medulla oblongata		inter-ventricular septum	appendix		bone marrow	ovary		colorectal adenocarcinoma SW480	fetal lung	human DNA 100 ng
H	pons	putamen		apex of the heart	colon ascending		trachea			lung carcinoma A549		human DNA 500 ng

Fig 4B

8/15



SUBSTITUTE SHEET (RULE 26)



SUBSTITUTE SHEET (RULE 26)

Fig 6 10/15




Fig 7 11/15




Fig 8 12/15




Fig 9 13/15




Fig 10 14/15





Fig 11 15/15



sequence_listing.ST25.txt
SEQUENCE LISTING

<110> Karolinska Innovations AB
Ingelman-Sundberg, Magnus
Karlgren, Maria
Gomez, Alvin

<120> Drug target in cancer therapy

<130> P05980PC00/HAM/em

<150> SE0203137-5

<151> 2002-10-24

<150> US 60/420,787

<151> 2002-10-24

<160> 10

<170> PatentIn version 3.1

<210> 1

<211> 27

<212> DNA

<213> Homo sapiens

<220>

<221> Primer

<222> (1)..(27)

<223>

<400> 1
gaggaggagg catcttcttc tcactctg

27

<210> 2

sequence_listing.ST25.txt

<211> 25

<212> DNA

<213> Homo sapiens

<220>

<221> primer

<222> (1)..(25)

<223>

<400> 2
ctggatcagg gcgtccacat agctg

25

<210> 3

<211> 23

<212> DNA

<213> homo sapiens

<220>

<221> primer

<222> (1)..(23)

<223>

<400> 3
agctatgtgg acgccctgat cca

23

<210> 4

<211> 23

<212> DNA

<213> homo sapiens

<220>

<221> primer

<222> (1)..(23)

<223>

<400> 4
acgcggtcta gctcctcctg cac

23

sequence_listing.ST25.txt

<210> 5
<211> 15
<212> PRT
<213> homo sapiens

<220>
<221> peptide
<222> (1)..(15)
<223>

<400> 5
Thr Met Arg Pro Arg Ala Gln Ala Leu Cys Ala Val Pro Arg Pro
1 5 10 15

<210> 6
<211> 27
<212> DNA
<213> homo sapiens

<220>
<221> primer
<222> (1)..(27)
<223>

<400> 6
gacagatcta tggccctgct gctcttg

27

<210> 7
<211> 28
<212> DNA
<213> homo sapiens

<220>
<221> primer
<222> (1)..(28)

sequence_listing.ST25.txt

<223>

<400> 7

gactctagac tagggcctgg gcaccgca

28

<210> 8

<211> 490

<212> PRT

<213> homo sapiens

<400> 8

Met Ala Leu Leu Leu Leu Phe Leu Gly Leu Leu Gly Leu Trp Gly
1 5 10 15Leu Leu Cys Ala Cys Ala Gln Asp Pro Ser Pro Ala Ala Arg Trp Pro
20 25 30Pro Gly Pro Arg Pro Leu Pro Leu Val Gly Asn Leu His Leu Leu Arg
35 40 45Leu Ser Gln Gln Asp Arg Ser Leu Met Glu Leu Ser Glu Arg Tyr Gly
50 55 60Pro Val Phe Thr Val His Leu Gly Arg Gln Lys Thr Val Val Leu Thr
65 70 75 80Gly Phe Glu Ala Val Lys Glu Ala Leu Ala Gly Pro Gly Gln Glu Leu
85 90 95Ala Asp Arg Pro Pro Ile Ala Ile Phe Gln Leu Ile Gln Arg Gly Gly
100 105 110Gly Ile Phe Phe Ser Ser Gly Ala Arg Trp Arg Ala Ala Arg Gln Phe
115 120 125Thr Val Arg Ala Leu His Ser Leu Gly Val Gly Arg Glu Pro Val Ala
130 135 140Asp Lys Ile Leu Gln Glu Leu Lys Cys Leu Ser Gly Gln Leu Asp Gly
145 150 155 160Tyr Arg Gly Arg Pro Phe Pro Leu Ala Leu Leu Gly Trp Ala Pro Ser
165 170 175Asn Ile Thr Phe Ala Leu Leu Phe Gly Arg Arg Phe Asp Tyr Arg Asp
180 185 190

sequence_listing.ST25.txt

Pro Val Phe Val Ser Leu Leu Gly Leu Ile Asp Glu Val Met Val Leu
 195 200 205
 Leu Gly Ser Pro Gly Leu Gln Leu Phe Asn Val Tyr Pro Trp Leu Gly
 210 215 220
 Ala Leu Leu Gln Leu His Arg Pro Val Leu Arg Lys Ile Glu Glu Val
 225 230 235 240
 Arg Ala Ile Leu Arg Thr Leu Leu Glu Ala Arg Arg Pro His Val Cys
 245 250 255
 Pro Gly Asp Pro Val Cys Ser Tyr Val Asp Ala Leu Ile Gln Gln Gly
 260 265 270
 Gln Gly Asp Asp Pro Glu Gly Leu Phe Ala Glu Ala Asn Ala Val Ala
 275 280 285
 Cys Thr Leu Asp Met Val Met Ala Gly Thr Glu Thr Thr Ser Ala Thr
 290 295 300
 Leu Gln Trp Ala Ala Leu Leu Met Gly Arg His Pro Asp Val Gln Gly
 305 310 315 320
 Arg Val Gln Glu Glu Leu Asp Arg Val Leu Gly Pro Gly Arg Thr Pro
 325 330 335
 Arg Leu Glu Asp Gln Gln Ala Leu Pro Tyr Thr Ser Ala Val Leu His
 340 345 350
 Glu Val Gln Arg Phe Ile Thr Leu Leu Pro His Val Pro Arg Cys Thr
 355 360 365
 Ala Ala Asp Thr Gln Leu Gly Gly Phe Leu Leu Pro Lys Gly Thr Pro
 370 375 380
 Val Ile Pro Leu Leu Thr Ser Val Leu Leu Asp Glu Thr Gln Trp Gln
 385 390 395 400
 Thr Pro Gly Gln Phe Asn Pro Gly His Phe Leu Asp Ala Asn Gly His
 405 410 415
 Phe Val Lys Arg Glu Ala Phe Leu Pro Phe Ser Ala Gly Arg Arg Val
 420 425 430
 Cys Val Gly Glu Arg Leu Ala Arg Thr Glu Leu Phe Leu Leu Phe Ala
 435 440 445
 Gly Leu Leu Gln Arg Tyr Arg Leu Leu Pro Pro Pro Gly Val Ser Pro
 450 455 460

sequence_listing.ST25.txt

Ala Ser Leu Asp Thr Thr Pro Ala Arg Ala Phe Thr Met Arg Pro Arg
 465 470 475 480

Ala Gln Ala Leu Cys Ala Val Pro Arg Pro
 485 490

<210> 9

<211> 2432

<212> DNA

<213> homo sapiens

<220>

<221> promoter

<222> (1)..(137)

<223>

<220>

<221> exon

<222> (138)..(1608)

<223>

<400> 9

```

gccctgcaaa ctgctcgttc cacattctcg ggggtggtggg gtgggtgggg tgcgggcacg      60
ccctcccgcg gaggcctata aggggtgcggg ggggacgggg cccaggaggg gagtggagcc      120
tcaccagcca cgtcctc  atg gcc ctg ctg ctc ttg ctg ttc ctg ggc ctc      170
                   Met Ala Leu Leu Leu Leu Leu Phe Leu Gly Leu
                   1             5             10

ctg ggg ctc tgg ggg ctg ctc tgc gcc tgc gcc caa gac ccc tcc cca      218
Leu Gly Leu Trp Gly Leu Leu Cys Ala Cys Ala Gln Asp Pro Ser Pro
                   15             20             25

gct gcc cgg tgg ccc ccg ggg cct cgc ccg ctg ccg ctc gtc ggg aac      266
Ala Ala Arg Trp Pro Pro Gly Pro Arg Pro Leu Pro Leu Val Gly Asn
                   30             35             40

ctg cac ttg ctg cgt ctg tcg caa cag gac cgg tcc ctg atg gag ctc      314
Leu His Leu Leu Arg Leu Ser Gln Gln Asp Arg Ser Leu Met Glu Leu
                   45             50             55

tca gaa cgc tac ggg ccg gtg ttc acc gtg cac ctg ggg cgc cag aag      362
Ser Glu Arg Tyr Gly Pro Val Phe Thr Val His Leu Gly Arg Gln Lys
60             65             70             75

acg gtg gtg ctg acg ggg ttc gag gcg gtc aaa gag gcg ctg gcg ggc      410
Thr Val Val Leu Thr Gly Phe Glu Ala Val Lys Glu Ala Leu Ala Gly
80             85             90

ccc ggg cag gag ctg gcc gac cgg cct ccc atc gcc atc ttc cag ctc      458
Sida 6

```


sequence_listing.ST25.txt

Pro	Gly	Gln	Glu 95	Leu	Ala	Asp	Arg	Pro 100	Pro	Ile	Ala	Ile 105	Phe	Gln	Leu	
atc	cag	cga	ggt	gga	ggc	atc	ttc	ttc	tca	tct	ggg	gcg	cgc	tgg	agg	506
Ile	Gln	Arg 110	Gly	Gly	Gly	Ile	Phe 115	Phe	Ser	Ser	Gly	Ala 120	Arg	Trp	Arg	
gct	gcc	cgc	cag	ttc	acg	gtg	cgt	gcc	ctg	cac	agc	ctg	ggc	gtg	ggc	554
Ala	Ala	Arg	Gln	Phe	Thr	Val 130	Arg	Ala	Leu	His	Ser 135	Leu	Gly	Val	Gly	
125																
cgg	gag	ccg	gtg	gct	gac	aag	att	ctg	cag	gag	ctg	aaa	tgc	ctc	tct	602
Arg	Glu	Pro	Val	Ala	Asp 145	Lys	Ile	Leu	Gln	Glu 150	Leu	Lys	Cys	Leu	Ser 155	
140																
ggg	cag	ctg	gat	ggc	tac	aga	ggc	cgg	ccc	ttc	ccg	ctg	gcc	cta	ctg	650
Gly	Gln	Leu	Asp	Gly 160	Tyr	Arg	Gly	Arg	Pro 165	Phe	Pro	Leu	Ala	Leu	Leu	
ggc	tgg	gct	ccc	tcc	aat	atc	acc	ttc	gcg	ctc	ctc	ttc	ggc	cgc	cga	698
Gly	Trp	Ala	Pro 175	Ser	Asn	Ile	Thr	Phe 180	Ala	Leu	Leu	Phe	Gly 185	Arg	Arg	
ttt	gac	tac	cgg	gac	ccc	gtg	ttt	gtg	tcc	ctg	ctg	ggt	ctc	atc	gat	746
Phe	Asp	Tyr 190	Arg	Asp	Pro	Val	Phe 195	Val	Ser	Leu	Leu	Gly 200	Leu	Ile	Asp	
gag	gtc	atg	gtc	ctc	ttg	ggg	tcc	cct	ggc	ctg	cag	ctg	ttc	aac	gtc	794
Glu	Val 205	Met	Val	Leu	Leu	Gly 210	Ser	Pro	Gly	Leu	Gln 215	Leu	Phe	Asn	Val	
tac	cca	tgg	ctc	ggg	gcc	ctg	ctc	cag	ctg	cac	cgg	ccc	gtc	ctg	cgc	842
Tyr	Pro	Trp	Leu	Gly	Ala 225	Leu	Leu	Gln	Leu	His 230	Arg	Pro	Val	Leu	Arg 235	
220																
aag	atc	gag	gag	gtc	cgt	gcc	att	ctg	agg	acc	ctc	ctg	gag	gcg	cgg	890
Lys	Ile	Glu	Glu	Val 240	Arg	Ala	Ile	Leu	Arg 245	Thr	Leu	Leu	Glu	Ala 250	Arg	
agg	ccc	cac	gtg	tgc	ccg	ggg	gac	ccc	gtg	tgc	agc	tat	gtg	gac	gcc	938
Arg	Pro	His	Val 255	Cys	Pro	Gly	Asp	Pro 260	Val	Cys	Ser	Tyr	Val 265	Asp	Ala	
ctg	atc	cag	cag	gga	cag	ggg	gat	gac	ccc	gag	ggc	ctg	ttt	gct	gag	986
Leu	Ile	Gln	Gln	Gly	Gln	Gly	Asp 275	Asp	Pro	Glu	Gly	Leu	Phe	Ala	Glu	
gcc	aac	gcg	gtg	gcc	tgc	acc	ctg	gac	atg	gtc	atg	gcc	ggg	acg	gag	1034
Ala	Asn	Ala	Val	Ala	Cys	Thr 290	Leu	Asp	Met	Val	Met 295	Ala	Gly	Thr	Glu	
285																
acg	acc	tcg	gcc	acg	ctg	cag	tgg	gcc	gca	ctt	ctg	atg	ggc	cgg	cac	1082
Thr	Thr	Ser	Ala	Thr	Leu 305	Gln	Trp	Ala	Ala	Leu 310	Leu	Met	Gly	Arg	His 315	
300																
ccg	gac	gtg	cag	ggc	cgg	gtg	cag	gag	gag	cta	gac	cgc	gtg	ctg	ggc	1130
Pro	Asp	Val	Gln	Gly 320	Arg	Val	Gln	Glu	Glu 325	Leu	Asp	Arg	Val	Leu 330	Gly	
cct	ggg	cgg	act	ccc	cgg	ctg	gag	gac	cag	cag	gct	ctg	ccc	tac	aca	1178
Pro	Gly	Arg	Thr 335	Pro	Arg	Leu	Glu	Asp 340	Gln	Gln	Ala	Leu	Pro 345	Tyr	Thr	
agc	gcc	gtg	ctc	cac	gag	gtg	cag	cgg	ttc	atc	acg	ctc	ctg	ccg	cac	1226
Ser	Ala	Val 350	Leu	His	Glu	Val	Gln 355	Arg	Phe	Ile	Thr	Leu 360	Leu	Pro	His	
gtg	ccc	cgc	tgc	acc	gcg	gcc	gac	aca	cag	ctg	ggc	ggc	ttc	ctg	ctc	1274

sequence_listing.ST25.txt

Val	Pro	Arg	Cys	Thr	Ala	Ala	Asp	Thr	Gln	Leu	Gly	Gly	Phe	Leu	Leu	
365						370					375					
ccc	aag	ggc	acg	ccc	gtg	att	ccc	ctg	ctg	acc	tcg	gtg	ctc	ctg	gat	1322
Pro	Lys	Gly	Thr	Pro	Val	Ile	Pro	Leu	Leu	Thr	Ser	Val	Leu	Leu	Asp	
380					385					390					395	
gag	aca	cag	tgg	cag	acc	cca	ggc	cag	ttc	aac	ccc	ggc	cat	ttc	ctg	1370
Glu	Thr	Gln	Trp	Gln	Thr	Pro	Gly	Gln	Phe	Asn	Pro	Gly	His	Phe	Leu	
				400				405						410		
gac	gcg	aat	ggg	cac	ttt	gtg	aag	cgg	gag	gcc	ttc	ctg	cct	ttc	tct	1418
Asp	Ala	Asn	Gly	His	Phe	Val	Lys	Arg	Glu	Ala	Phe	Leu	Pro	Phe	Ser	
			415					420					425			
gca	ggc	cgc	cgc	gtc	tgt	gtt	ggg	gag	cgc	ctg	gcc	agg	acc	gag	ctc	1466
Ala	Gly	Arg	Arg	Val	Cys	Val	Gly	Glu	Arg	Leu	Ala	Arg	Thr	Glu	Leu	
		430					435					440				
ttc	ctg	ctg	ttt	gcc	ggc	ctc	ctg	cag	agg	tac	cgc	ctg	ctg	ccc	ccg	1514
Phe	Leu	Leu	Phe	Ala	Gly	Leu	Leu	Gln	Arg	Tyr	Arg	Leu	Leu	Pro	Pro	
	445					450					455					
cct	ggc	gtc	agt	ccg	gcc	tcc	ctg	gac	acc	acg	ccc	gcc	cgg	gct	ttt	1562
Pro	Gly	Val	Ser	Pro	Ala	Ser	Leu	Asp	Thr	Thr	Pro	Ala	Arg	Ala	Phe	
460					465					470					475	
acc	atg	agg	ccg	agg	gcc	cag	gcc	ctg	tgt	gcg	gtg	ccc	agg	ccc	t	1608
Thr	Met	Arg	Pro	Arg	Ala	Gln	Ala	Leu	Cys	Ala	Val	Pro	Arg	Pro		
				480					485					490		
aggagctccc	ccagccccc	ggctctctg	accactcccc	tcccagccct	gggtcctccc											1668
accctctctc	ctcccacccc	acagctcgga	ctgctctggg	agggccctga	ggactccac											1728
cctcaccccc	acccccacag	ggtcagcaac	tgcttccggt	tacaccagc	actaccctg											1788
cccgaccctg	tgggaccccc	accctctga	tgctgtctgc	agctcagtcc	ctgccagccc											1848
ccaggagcgc	ctccagggcc	ccgcccactc	tcccaccct	gaagctgcac	tcccaccac											1908
ctagctcccc	ccagggccccc	ccagcaccta	cagctggggc	tgcagggaga	caacgggtgg											1968
ctgcatccag	ccagagacag	gcgcaggtgg	gtgtcctcag	cgtgcgagcc	ctgcaccccc											2028
caggctcctgg	gactcctgca	gacccactc	cattcccgt	cctggaacac	ttcctgcagc											2088
tgtgcctgga	ggcagtcggc	ctgcagtgcc	agactctgag	ccaagccact	ggggccatgc											2148
gtatgactgg	tgcagggagg	caaggccac	attctccttc	agagacaggc	actggcgcca											2208
gaggcttcct	tggggcgggg	ggagggcacc	tcagcccctg	aagacaagca	gcactgcagt											2268
ggcaaaaatg	gaaacactga	cccgggtcgg	tggctcatgc	ctgtaatccc	agcactttgg											2328
gaggccgagg	caggcgaatc	acgaggtcag	gagttcgaga	ccagcctgcc	caacatggtg											2388
aaaccctgtc	tctactaaaa	atacaaaaaa	attaaccgag	catg												2432

<210> 10

<211> 137

<212> DNA

<213> Homo sapiens

sequence_listing.ST25.txt

<220>

<221> promoter

<222> (1)..(137)

<223> -137 to -1 5'UTR of CYP2W1 gene

<400> 10

gccctgcaaa ctgctcggtc cacattctcg ggggtggtggg gtgggtgggg tgcgggcacg 60

ccctcccgcg gaggcctata aggggtgcggg ggggacgggg cccaggaggg gagtggagcc 120

tcaccagcca cgtcctc 137

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 2003/001652

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: A61K 38/44, G01N 33/574

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: A61K, G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL (Online) Accession no. Q8TAV3, 1 June 2002, retrieved from EBI, 100% identity in 434 amino acids overlap with SEQ ID No 8 --	1-10
A	YOUSSEF JOUNAIDI et al "Frequent, Moderate-Dose Cyclophosphamide Administration Improves the Efficacy of Cytochrome P-450/Cytochrome P-450 Reductasebased Cancer Gene Therapy1", Cancer Research, June 1 2001, Vol. 61, p. 4437- 4444., --	1-7, 10

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

28 January 2004

Date of mailing of the international search report

04-02-2004

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Yvonne Siösteen/ELY

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE 2003/001652

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	STN International, File CAPLUS, CAPLUS accession no. 1997:210366, document no. 126:221319, Oyama, Tsunehiro et al: "Cytochrome P450 2E1 polymorphism as a risk factor for lung cancer: in relation to p53 gene mutation"; & Anticancer Research (1997), 17(1B), 583-587 --	1-7,10
A	BIOSIS, accession no. PREV200200495945, Itoga Sakae et al: "Tandem repeat polymorphism of the CYP2E1 gene: An association study with esophageal cancer and lung cancer", Alcoholism Clinical and Experimental Research, August, 2002, Vol. 26, No. 8 Supplement, pp. 15S-19S --	1-7,10
A	WO 0158455 A1 (PHARMACIA & UPJOHN S.P.A.), 16 August 2001 (16.08.2001), see claim 29 --	1-7,10
A	STN International, File CAPLUS, CAPLUS accession no. 1996:580855, document no. 125:292401, Vasquez, Hernan G. et al: "Effect of cytochrome P450 inducers on expression of oncogenes and a tumor suppressor gene in human colon cancer cells"; & International Journal of Oncology (1996), 9(3), pp. 427-431 --	1-7,10
A	MAGNUS INGELMAN-SUNDBERG, "Genetic susceptibility to adverse effects of drugs and environmental toxicants The role of the CYP family of enzymes", Mutation Research 2001, Vol. 482, pp. 11-19 see page 13 table 1 --	1-7,10
A	DANIEL W NEBERT et al: "Clinical importance of the cytochromes P450", THE LANCET, 12 October 2002, Vol. 360, pp. 1155-62, see page 1157, line 13-15 -- -----	1-7,10

INTERNATIONAL SEARCH REPORT
Information on patent family members

24/12/2003

International application No.

PCT/SE 2003/001652

WO	0158455	A1	16/08/2001	AU	3022901	A	20/08/2001
				AU	6707900	A	19/03/2001
				BR	0108283	A	29/10/2002
				CA	2398840	A	16/08/2001
				CN	1398185	T	19/02/2003
				EE	200200440	A	15/12/2003
				EP	1208642	A	29/05/2002
				EP	1267889	A	02/01/2003
				GB	0003201	D	00/00/0000
				JP	2003524645	T	19/08/2003
				US	2003153539	A	14/08/2003

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.